



THE UNIVERSITY of EDINBURGH

Edinburgh Research Explorer

New loci associated with kidney function and chronic kidney disease

Citation for published version:

Köttgen, A, Pattaro, C, Böger, CA, Fuchsberger, C, Olden, M, Glazer, NL, Parsa, A, Gao, X, Yang, Q, Smith, AV, O'Connell, JR, Li, M, Schmidt, H, Tanaka, T, Isaacs, A, Ketkar, S, Hwang, S-J, Johnson, AD, Dehghan, A, Teumer, A, Paré, G, Atkinson, EJ, Zeller, T, Lohman, K, Cornelis, MC, Probst-Hensch, NM, Kronenberg, F, Tönjes, A, Hayward, C, Aspelund, T, Eiriksdottir, G, Launer, LJ, Harris, TB, Rasmussen, E, Mitchell, BD, Arking, DE, Boerwinkle, E, Struchalin, M, Cavalieri, M, Singleton, A, Giallauria, F, Metter, J, de Boer, IH, Haritunians, T, Lumley, T, Siscovick, D, Psaty, BM, Zillikens, MC, Oostra, BA, Feitosa, M, Province, M, de Andrade, M, Turner, ST, Schillert, A, Ziegler, A, Wild, PS, Schnabel, RB, Wilde, S, Munzel, TF, Leak, TS, Illig, T, Klopp, N, Meisinger, C, Wichmann, H-E, Koenig, W, Zgaga, L, Zemunik, T, Kolcic, I, Minelli, C, Hu, FB, Johansson, A, Igl, W, Zaboli, G, Wild, SH, Wright, AF, Campbell, H, Ellinghaus, D, Schreiber, S, Aulchenko, YS, Felix, JF, Rivadeneira, F, Uitterlinden, AG, Hofman, A, Imboden, M, Nitsch, D, Brandstätter, A, Kollerits, B, Kedenko, L, Mägi, R, Stumvoll, M, Kovacs, P, Boban, M, Campbell, S, Endlich, K, Völzke, H, Kroemer, HK, Nauck, M, Völker, U, Polasek, O, Vitart, V, Badola, S, Parker, AN, Ridker, PM, Kardia, SLR, Blankenberg, S, Liu, Y, Curhan, GC, Franke, A, Roach, T, Paulweber, B, Prokopenko, I, Wang, W, Gudnason, V, Shuldiner, AR, Coresh, J, Schmidt, R, Ferrucci, L, Shlipak, MG, van Duijn, CM, Borecki, I, Krämer, BK, Rudan, I, Gyllenstein, U, Wilson, JF, Witteman, JC, Pramstaller, PP, Rettig, R, Hastie, N, Chasman, DI, Kao, WH, Heid, IM & Fox, CS 2010, 'New loci associated with kidney function and chronic kidney disease', *Nature Genetics*, vol. 42, no. 5, pp. 376-384. <https://doi.org/10.1038/ng.568>

Digital Object Identifier (DOI):

[10.1038/ng.568](https://doi.org/10.1038/ng.568)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Nature Genetics

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Download date: 04. May. 2023



Published in final edited form as:

Nat Genet. 2010 May ; 42(5): 376–384. doi:10.1038/ng.568.

Multiple New Loci Associated with Kidney Function and Chronic Kidney Disease: The CKDGen consortium

Anna Köttgen, MD MPH^{*,†,1}, Cristian Pattaro, PhD^{*,†,2}, Carsten A. Böger, MD^{+,*,3}, Christian Fuchsberger, PhD^{+,*,2}, Matthias Olden, PhD⁴, Nicole L. Glazer, PhD MPH^{+,5}, Afshin Parsa, MD MPH^{+,6}, Xiaoyi Gao, PhD^{+,7}, Qiong Yang, PhD⁸, Albert V. Smith, PhD⁹, Jeffrey R. O'Connell, PhD⁶, Man Li, MS¹, Helena Schmidt, MD PhD¹⁰, Toshiko Tanaka, PhD¹¹, Aaron Isaacs, PhD¹², Shamika Ketkar, MS⁷, Shih-Jen Hwang, PhD¹³, Andrew D. Johnson, PhD¹³, Abbas Dehghan, MD MSc¹⁴, Alexander Teumer, MSc¹⁵, Guillaume Paré, MD MSc¹⁶, Elizabeth J. Atkinson, MS¹⁷, Tanja Zeller, PhD¹⁸, Kurt Lohman, MSTAT¹⁹, Marilyn C. Cornelis, PhD²⁰, Nicole M. Probst-Hensch, PhD MPH²¹, Florian Kronenberg, MD²², Anke Tönjes, MD²³, Caroline Hayward, PhD²⁴, Thor Aspelund, PhD^{9,25}, Gudny Eiriksdottir, MSc⁹, Lenore Launer, PhD²⁶, Tamara B. Harris, MD MPH²⁶, Evadnie Rappersaud, PhD²⁷, Braxton D. Mitchell, PhD⁶, Eric Boerwinkle, PhD²⁸, Maksim Struchalin, PhD²⁹, Margherita Cavalieri, MD³⁰, Andrew Singleton, PhD³¹, Francesco Giallauria, MD PhD³², Jeffery Metter, PhD³², Ian de Boer, MD MS³³, Talin Haritunians, PhD³⁴, Thomas Lumley, PhD³⁵, David Siscovick, MD MPH³⁶, Bruce M. Psaty, MD PhD³⁷, M. Carola Zillikens, MD¹², Ben A. Oostra, PhD¹², Mary Feitosa, PhD⁷, Michael Province, PhD⁷, Daniel Levy, MD¹³, Mariza de Andrade, PhD¹⁷, Stephen T. Turner, MD³⁸, Arne Schillert, PhD³⁹, Andreas Ziegler, PhD³⁹, Philipp S. Wild, MD¹⁸, Renate B. Schnabel, MD MSc¹⁸, Sandra Wilde, BA¹⁸, Thomas F. Muenzel, MD¹⁸, Tennille S Leak, PhD⁴⁰, Thomas Illig, PhD⁴¹, Norman Klopp, PhD⁴¹, Christa Meisinger, MD MPH⁴¹, H.-Erich Wichmann, MD⁴², Wolfgang Koenig, MD⁴³, Lina Zgaga, MD⁴⁴, Tatijana Zemunik, MD PhD⁴⁵, Ivana Kolcic, MD PhD⁴⁶, Cosetta Minelli, MD PhD², Frank B. Hu, MD PhD²⁰, Åsa Johansson, PhD⁴⁷, Wilmar Igl, PhD⁴⁷, Ghazal Zabolli, PhD⁴⁷, Sarah H Wild, PhD⁴⁸, Alan F Wright, PhD⁴⁹, Harry Campbell, MD⁴⁸, David Ellinghaus, MSc⁵⁰, Stefan Schreiber, MD⁵⁰, Yuri S Aulchenko, PhD⁵¹, Fernando Rivadeneira, MD PhD⁵¹, Andre G Uitterlinden, PhD⁵¹, Albert Hofman, MD PhD⁵¹, Medea Imboden, PhD²¹, Dorothea Nitsch, MD MSc⁵², Anita Brandstätter, PhD²², Barbara Kollerits, PhD MPH²², Lyudmyla Kedenko, PhD⁵³, Reedik Mägi, PhD⁵⁴, Michael Stumvoll, MD²³, Peter Kovacs, PhD²³, Mladen Boban, MD PhD⁵⁵, Susan Campbell, BSc⁵⁶, Karlhans Endlich, MD⁵⁷, Henry Völzke, MD⁵⁸, Heyo K. Kroemer, PhD⁵⁹, Matthias Nauck, MD⁶⁰, Uwe Völker, PhD⁶¹, Ozren Polasek, MD PhD⁴⁶, Veronique Vitart, PhD²⁴, Sunita Badola, MS⁶², Alexander N. Parker, PhD⁶², Paul M. Ridker, MD MPH⁶³, Sharon L. R. Kardia, PhD⁶⁴, Stefan Blankenberg, MD¹⁸, Yongmei Liu, MD PhD⁶⁵, Gary C. Curhan, MD ScD⁶⁶, Andre Franke, PhD⁵⁰, Thierry Rochat, MD⁶⁷, Bernhard Paulweber, MD⁵³, Inga Prokopenko, PhD⁵⁴, Wei Wang, MD PhD⁶⁸, Vilundur Gudnason, MD PhD^{25,9}, Alan R. Shuldiner, MD⁶, Josef Coresh, MD PhD⁶⁹, Reinhold Schmidt, MD³⁰, Luigi Ferrucci, MD PhD³², Michael G. Shlipak, MD MPH⁷⁰, Cornelia M. van Duijn, PhD¹², Ingrid Borecki, PhD⁷, Bernhard K. Krämer, MD⁷¹, Igor Rudan, MD PhD⁴⁴, Ulf Gyllenstein,

Corresponding Authors: Caroline S. Fox MD MPH, NHLBI's Framingham Heart Study, 73 Mt Wayte Ave Suite #2, Framingham MA 01702, foxca@nhlbi.nih.gov, Iris M. Heid PhD, Department of Epidemiology and Preventive Medicine, Regensburg University Medical Center Franz-Josef-Strauß-Allee 11, 93053 Regensburg, Germany, iris.heid@klinik.uni-regensburg.de, W. H. Kao PhD, Department of Epidemiology, Welch Center for Prevention, Epidemiology, and Clinical Research, Johns Hopkins University, 615 N. Wolfe St., Room 6513, Baltimore, MD 21205, wkao@jhsph.edu, Daniel I. Chasman PhD, Brigham and Women's Hospital, 900 Commonwealth Avenue, East Boston, MA 02215, dchasman@rics.bwh.harvard.edu.

*These co-authors contributed equally to this work

**These senior authors jointly oversaw this work

+Writing Group

PhD⁴⁷, James F. Wilson, DPhil⁴⁸, Jacqueline C. Witteman, PhD⁵¹, Peter P. Pramstaller, MD², Rainer Rettig, MD⁷², Nick Hastie, PhD²⁴, Daniel I. Chasman, PhD⁶³, W. H. Kao, PhD⁶⁹, Iris M. Heid, PhD⁷³, and Caroline S. Fox, MD MPH⁷⁴

¹Department of Epidemiology, Johns Hopkins University, 615 N. Wolfe St., Room 6021, Baltimore, MD 21205. ²Institute of Genetic Medicine, European Academy Bozen/Bolzano (EURAC), Bolzano, Italy and Affiliated Institute of the University of Lübeck, Germany, Viale Druso, 1 - 39100 Bolzano (Italy). ³Department of Internal Medicine II, University Medical Center Regensburg, Franz-Josef-Strauss-Allee 11, 93042 Regensburg, Germany. ⁴Department of Internal Medicine II, University Medical Center Regensburg and the Department of Epidemiology and Preventive Medicine, University Medical Center Regensburg, Franz-Josef-Strauss-Allee 11, 93042 Regensburg, Germany. ⁵Cardiovascular Health Research Unit and Department of Medicine, University of Washington, Seattle, WA, 1730 Minor Ave, Suite 1360, Seattle, WA, 98101. ⁶University of Maryland Medical School, 660 W Redwood St, RM 494, Baltimore, MD, 21212. ⁷Division of Statistical Genomics; Washington University School of Medicine, 4444 Forest Park Blvd. - Suite 8506; St Louis, MO 63108. ⁸Department of Biostatistics, Boston University School of Public Health, 715 Albany Street, Boston, MA 02118. ⁹Icelandic Heart Association, Hjartavernd, Holtasmara 1, 201 Kopavogur, Iceland. ¹⁰Austrian Stroke Prevention Study, Institute of Molecular Biology and Biochemistry and University Clinic of Neurology, Department of Neurogeriatrics, Medical University Graz, Auenbruggerplatz 22, 8036, Austria. ¹¹Clinical Research Branch, National Institute on Aging and Medstar Research Institute, Baltimore MD 21250, USA. ¹²Genetic Epidemiology Unit, Department of Epidemiology, Erasmus University Medical Center, PO Box 2040, 3000CA Rotterdam, the Netherlands. ¹³NHLBI's Framingham Heart Study and the Center for Population Studies, 73 Mt Wayte Ave Suite #2, Framingham MA 01702. ¹⁴Department of Epidemiology, Erasmus Medical Center, P.O. Box 2040, 3000 CA, Rotterdam, The Netherlands. ¹⁵Interfaculty Institute for Genetics and Functional Genomics, Ernst-Moritz-Arndt-University Greifswald, 17487 Greifswald, Germany. ¹⁶Department of Pathology & Molecular Medicine, McMaster University, 1200 Main St. W MDCL Rm. 3206, Hamilton, ON Canada, L8N3Z5. ¹⁷Division of Biomedical Statistics and Informatics, Mayo Clinic, 200 First Street SW, Rochester, MN 55905. ¹⁸Department of Medicine II, University Medical Center Mainz, Langenbeckstr. 1, 55131 Mainz, Germany. ¹⁹Department of Biostatistical Sciences, Wake Forest University, Division of Public Health Sciences, Medical Center Blvd, Winston-Salem, NC 27157. ²⁰Harvard School of Public Health Department of Nutrition, 665 Huntington Avenue, Building 2, Boston MA, 02115. ²¹Institute of Social and Preventive Medicine (ISPM) at Swiss Tropical Institute Basel STI - An Associated Institute of the University of Basel, Steinengraben 49, 4051 Basel, Switzerland. ²²Innsbruck Medical University, Division of Genetic Epidemiology, Schoepfstraße 41, 6020 Innsbruck, Austria. ²³Department of Medicine, University of Leipzig, Liebigstr. 18, 04103 Leipzig, Germany. ²⁴MRC Human Genetics Unit, Institute of Genetics and Molecular Medicine, Western General Hospital, Crewe Road, Edinburgh EH4 2XU, Scotland. ²⁵University of Iceland, Reykjavik, Iceland. ²⁶Laboratory of Epidemiology, Demography, and Biometry, NIA, Gateway Building, 3C309, 7201 Wisconsin Ave, Bethesda MD 20892-9205. ²⁷Miami Institute for Human Genomics, 1501 NW 10th ave, Miami, FL 33136. ²⁸Human Genetics Center, University of Texas Health Science Center, Houston, TX, 1200 Herman Pressler Drive, Houston, TX 77030. ²⁹Department of Epidemiology and Biostatistics, Department of Forensic Molecular Biology, Dr. Molewaterplein 50-603015 GE Rotterdam; The Netherlands. ³⁰Austrian Stroke Prevention Study, University Clinic of Neurology, Department of Neurogeriatrics, Medical University Graz, Auenbruggerplatz 22, 8036 Graz, Austria. ³¹Laboratory of Neurogenetics, National Institute on Aging, Bethesda MD, USA. ³²Clinical Research Branch, National Institute on Aging, Baltimore MD 21250, USA. ³³Division of Nephrology, University of Washington, Box 357183, 1959 NE Pacific St, Seattle, WA 98195. ³⁴Medical Genetics Institute, Cedars-Sinai Medical Center, Los Angeles, CA, USA, Cedars-Sinai Medical Center, Davis Building Room 4094B, 110 N. George Burns Road, Los Angeles, CA 90048. ³⁵Department of Biostatistics,

University of Washington, Seattle, WA, USA, 1730 Minor Ave, Suite 1360, Seattle, WA, 98101.

³⁶Departments of Epidemiology and Medicine, University of Washington, Seattle, WA, USA, 1730 Minor Ave, Suite 1360, Seattle, WA, 98101. ³⁷Cardiovascular Health Research Unit, Departments of Medicine, Epidemiology, and Health Services, University of Washington and Group Health Research Institute, Group Health, 1730 Minor Ave, Suite 1360, Seattle, WA, 98101.

³⁸Dept. of Internal Medicine, Division of Nephrology and Hypertension, Mayo Clinic, 200 First Street SW, Rochester, MN 55905. ³⁹Institute for Medical Biometry and Statistics, University at Luebeck, Maria-Goeppert-Str. 1, 23562 Luebeck, Germany. ⁴⁰Department of Epidemiology, Graduate School of Public Health, University of Pittsburgh, Graduate School of Public Health, 130 N. Bellefield Avenue, Room 512, Pittsburgh PA 15213. ⁴¹Institute of Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Ingolstädter Landstr. 1, 85764 Neuherberg, Germany. ⁴²Institute of Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Institute of Medical Informatics, Biometry and Epidemiology, Ludwig-Maximilians-Universität and Klinikum Grosshadern, Ingolstädter Landstr. 1, 85764 Neuherberg, Germany. ⁴³Zentrum für Innere Medizin, Klinik für Innere Medizin II - Kardiologie, Universitätsklinikum Ulm, Albert-Einstein-Allee 23, 89081 Ulm, Germany. ⁴⁴Center for Population Health Sciences, the University of Edinburgh Medical School, UK, Teviot Place, Edinburgh, EH8 9AG, Scotland. ⁴⁵Croatian Centre for Global Health, University of Split Medical School, Croatia, Croatian Centre for Global Health, University of Split Medical School, Split, Croatia. ⁴⁶Gen-Info Ltd, Ruzmarinka 17, 10000 Zagreb, Croatia. ⁴⁷Uppsala University, Genetics and Pathology, Rudbeck Laboratory, Uppsala University, SE-751 85, Uppsala, Sweden. ⁴⁸Centre for Population Health Sciences, University of Edinburgh, Teviot Place, Edinburgh, EH8 9AG, Scotland. ⁴⁹MRC Human Genetics Unit, Institute of Genetics and Molecular Medicine, Western General Hospital, Crewe Road, Edinburgh EH4 2XU, Scotland. ⁵⁰Institute of Clinical Molecular Biology, Christian-Albrechts University, Schittenhelmstr.12, 24105 Kiel, Germany. ⁵¹Department of Epidemiology, Erasmus Medical Center, P.O. Box 2040, 3000 CA, Rotterdam, The Netherlands. ⁵²Department of Epidemiology and Population Health, London School of Hygiene & Tropical Medicine, Keppel Street, London, United Kingdom. ⁵³First Department of Internal Medicine, Paracelsus Medical University, Salzburg, Austria, Müllner Hauptstr. 48, 5020 Salzburg, Austria. ⁵⁴Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, OX3 7BN, UK, Oxford Centre for Diabetes, Endocrinology and Metabolism, University of Oxford, UK, OX3 7LJ. ⁵⁵Croatian Centre for Global Health, University of Split Medical School, Croatia, Croatian Centre for Global Health, University of Split Medical School, Split, Croatia. ⁵⁶MRC Human Genetics Unit, Institute of Genetics and Molecular Medicine, Western General Hospital, Crewe Road, Edinburgh EH4 2XU, Scotland. ⁵⁷Institute of Anatomy and Cell Biology, Ernst-Moritz-Arndt-University Greifswald, 17487 Greifswald, Germany. ⁵⁸Institute for Community Medicine, Ernst-Moritz-Arndt-University Greifswald, 17487 Greifswald, Germany. ⁵⁹Institute of Pharmacology, Ernst-Moritz-Arndt-University Greifswald, 17487 Greifswald, Germany. ⁶⁰Institute of Clinical Chemistry and Laboratory Medicine, Ernst-Moritz-Arndt-University Greifswald, 17487 Greifswald, Germany. ⁶¹Interfaculty Institute for Genetics and Functional Genomics, Ernst-Moritz-Arndt-University Greifswald, Interfaculty Institute for Genetics and Functional Genomics, Ernst-Moritz-Arndt-University Greifswald, 17487. ⁶²Amgen, 360 Binney Street, Cambridge MA 02142. ⁶³Brigham and Women's Hospital, 900 Commonwealth Avenue, East, Boston, MA 02215. ⁶⁴University of Michigan School of Public Health, University of Michigan Department of Epidemiology, 109 Observatory, #4605, Ann Arbor, MI 48109-2029. ⁶⁵Department of Epidemiology and Prevention, Wake Forest University, Division of Public Health Sciences, Medical Center Blvd, Winston-Salem, NC 27157. ⁶⁶Brigham and Women's Hospital, Harvard Medical School, 181 Longwood Avenue, Boston, MA 02115. ⁶⁷University Hospitals of Geneva, 24, Rue Micheli-du-Crest, 1211 Geneve, 14, Switzerland. ⁶⁸School of Public Health and Family Medicine, Capital Medical University, Beijing, China, Croatian Centre for Global Health, University of Split Medical School, Split, Croatia. ⁶⁹Welch Center for Prevention, Epidemiology & Clinical

Research, Johns Hopkins University, 2024 E. Monument St., Baltimore, MD 21287, USA. ⁷⁰General Internal Medicine, University of California, San Francisco, 4150 Clement St., San Francisco, CA 94121. ⁷¹Department of Medicine I. Marien Hospital Herne, Medical Center of the Ruhr Universität Bochum, Hölkeskampring 40. 44625 Herne, Germany. ⁷²Institute of Physiology, Ernst-Moritz-Arndt-University Greifswald, 17487 Greifswald, Germany. ⁷³Department of Epidemiology and Preventive Medicine, University Medical Center Regensburg and the Institute of Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Franz-Josef-Strauss-Allee 11, 93042 Regensburg, Germany. ⁷⁴NHLBI's Framingham Heart Study, Center for Population Studies, and Harvard Medical School, 73 Mt Wayte Ave Suite #2, Framingham MA 01702.

Abstract

Chronic kidney disease (CKD) is a significant public health problem, and recent genetic studies have identified common CKD susceptibility variants. The CKDGen consortium performed a meta-analysis of genome-wide association data in 67,093 Caucasian individuals from 20 population-based studies to identify new susceptibility loci for reduced renal function, estimated by serum creatinine (eGFR_{crea}), cystatin C (eGFR_{cys}), and CKD (eGFR_{crea} <60 ml/min/1.73m²; n = 5,807 CKD cases). Follow-up of the 23 genome-wide significant loci ($p < 5 \times 10^{-8}$) in 22,982 replication samples identified 13 novel loci for renal function and CKD (in or near *LASS2*, *GCKR*, *ALMS1*, *TFDP2*, *DAB2*, *SLC34A1*, *VEGFA*, *PRKAG2*, *PIP5K1B*, *ATXN2*, *DACH1*, *UBE2Q2*, and *SLC7A9*) and 7 creatinine production and secretion loci (*CPS1*, *SLC22A2*, *TMEM60*, *WDR37*, *SLC6A13*, *WDR72*, *BCAS3*). These results further our understanding of biologic mechanisms of kidney function by identifying loci potentially influencing nephrogenesis, podocyte function, angiogenesis, solute transport, and metabolic functions of the kidney.

Keywords

genome-wide association; renal disease; population-based; genetics; chronic kidney disease

Introduction

Chronic kidney disease (CKD) is estimated to affect over 13% of adults¹ and is increasing in prevalence.^{1,2} This poses a significant global disease burden as the risk for end stage renal disease (ESRD), cardiovascular morbidity, and mortality increases with declining glomerular filtration rate (GFR),³ the most commonly used measure of kidney function. In addition, CKD incurs substantial expenditures in the US,⁴ with similar trends expected globally.⁵

Despite the increasing prevalence of CKD, our understanding of the underlying risk factors and pathophysiologic mechanisms remains incomplete.⁵ Hypertension and diabetes are major risk factors for CKD.⁶ However, the marked variability in the development of CKD in the setting of hypertension and diabetes demonstrates that additional underlying factors contribute to its etiology.⁷ In particular, studies have consistently demonstrated important genetic contributions to estimated GFR (eGFR), CKD and ESRD.^{8,9} Using genome-wide association, we have recently identified susceptibility variants for renal function and CKD at the *UMOD*, *SHROOM3*, and *STC1* loci in nearly 20,000 individuals.¹⁰ Together, single nucleotide polymorphisms (SNPs) at these loci explain only 0.43% of the variance in eGFR,¹⁰ suggesting that additional loci remain to be identified.

Thus, we have now performed a genome-wide association meta-analysis in 67,093 Caucasian participants from 20 general population-based cohorts within the CKDGen consortium, followed by independent replication of our findings in 22,982 Caucasian individuals. We analyzed GFR estimated from serum creatinine by the Modification of Diet in Renal Disease (MDRD) Study equation (eGFR_{crea}) as well as CKD (eGFR_{crea} <60 ml/min/1.73m²). To discriminate true susceptibility loci for renal function from those related to creatinine production and secretion, we used GFR estimated from a second serum marker of kidney function, cystatin C (eGFR_{cys}).

RESULTS

Study Samples

Overall, 90,075 individuals (67,093 in Stage 1 discovery and 22,982 Stage 2 replication) contributed information to the analysis of eGFR_{crea}, 84,740 individuals (62,237 Stage 1 discovery and 22,503 Stage 2 replication) to the analysis of CKD, and 26,071 to the analysis of eGFR_{cys} (20,957 Stage 1 discovery and 5,114 Stage 2 replication; Table 1).

Meta-Analysis in CKDGen Stage 1 Discovery Cohorts

Table 2 summarizes information for the 28 genomic loci that contained at least one genome-wide significant SNP association ($p < 5 \times 10^{-8}$) for any of the three discovery traits; the SNP with the lowest p-value at each locus is presented. In addition to confirming 5 known loci,¹⁰ we identified 23 novel loci containing genome-wide significant SNPs (p-values between 4.5×10^{-8} and 3.8×10^{-12}): 20 SNPs were identified in association with eGFR_{crea}, 2 SNPs with CKD, and 1 SNP with eGFR_{cys}. Of note, rs7805747 in the *PRKAG2* gene was identified in discovery analyses for both eGFR_{crea} and CKD, as was the known lead SNP at the *UMOD* locus.

Figure 1A shows the genome-wide $-\log_{10}$ p-value plot from Stage 1 discovery association analyses with eGFR_{crea}, Figure 1B with CKD, and Figure 1C shows the eGFR_{cys} results. The respective quantile-quantile plots are presented in Supplementary Figure 1. Study-specific and median imputation quality for the lead SNPs can be found in Supplementary Table 2.

Corroborating Evidence with eGFR_{cys}

Because serum creatinine concentration is influenced both by renal function as well as by creatinine production or secretion, we used eGFR_{cys} as a second measure of renal function to help distinguish between true renal function loci and creatinine production or secretion loci. Thus, of the 23 newly discovered loci, 16 were classified as renal function loci based on a direction-consistent association with eGFR_{cys} with p-value of <0.05 (Table 2 and Supplementary Table 3), and 7 were classified as loci related to creatinine metabolism. One SNP, rs653178 at the *ATXN2* locus, was identified primarily in association with eGFR_{cys}.

Stage 2: Independent Replication of Genome-wide Significant Discovery Findings

The lead SNP at each of the 20 novel loci for eGFR_{crea}, the 2 loci for CKD, and the novel locus for eGFR_{cys} were evaluated for independent replication in Stage 2 analyses with the respective discovery trait. After meta-analysis of Stage 1 discovery and Stage 2 replication results, the associations for all but 3 SNPs (rs16864170 at *SOX11*, rs1933182 at *SYPL2*, rs4014195 at *RNASEH2C*) became more significant (Table 2). Additionally, 16 of these 20 SNPs also showed a significant association in the replication samples alone (one-sided p-value <0.0025 for eGFR_{crea}, p <0.025 for CKD, and p <0.05 for eGFR_{cys}, Table 2). Thus, after integrating evidence for replication with association results for eGFR_{cys}, 13 replicated loci for renal function and 7 replicated loci likely related to creatinine metabolism were

identified. Of note, rs653178 at the *ATXN2* locus, which was identified primarily in association with eGFRcys, was also associated with eGFRcrea in the combined discovery and replication results. Regional association plots for all replicated loci related to renal function are shown in Supplementary Figure 2.

Of the SNPs that were validated by Stage 2 replication and were also associated with eGFRcys, *ALMS1*, *DAB2*, *SLC34A1*, *PRKAG2*, *VEGFA*, *DACH1*, and *SLC7A9* can be linked to renal function and/or disease and are highlighted together with *LASS2* and *GCKR* in Box 1. The remaining novel renal function loci were located in or close to *TFDP2*, *PIP5K1B*, *UBE2Q2*, and *ATXN2*. In spite of a lack of clear biological connection to renal function at these loci, findings were consistent across eGFRcrea, CKD, and eGFRcys analyses. More information on these genes is presented in Supplementary Box 1.

Of the 20 SNPs that replicated, the remaining 7 SNPs were not associated with eGFRcys and hence were considered as likely creatinine production or secretion loci (*CPS1*, *SLC22A2*, *TMEM60*, *WDR37*, *SLC6A13*, *WDR72*, and *BCAS3*). More information on these loci is presented in Supplementary Box 1.

SNPs associated with eGFRcrea are also associated with CKD

The majority of the 13 validated renal function loci were nominally associated with CKD (Table 3), underscoring how the use of intermediate phenotypes can provide insight into disease-based traits. The odds ratios associated with CKD for each additional copy of the minor allele ranged from 0.93 to 1.19, and minor allele frequencies ranged from 0.20 to 0.50.

Stratified Analyses and Measures of Clinical Relevance

Since diabetes mellitus and hypertension are major risk factors for kidney disease, we investigated the association of the replicated renal function loci with eGFRcrea stratified by diabetes or hypertension status in the discovery cohorts. None of the SNPs reported in Table 2 differed significantly across strata of diabetes and hypertension ($p < 0.008$, Bonferroni-corrected alpha of 0.1 for 13 tests).

The 13 confirmed and the three previously identified renal function loci account for 1.4% of the variation in eGFRcrea. A genetic risk score was computed using all 16 validated renal loci (13 novel, 3 known) and data from the ARIC study. Across categories of the genetic risk score, mean eGFRcrea ranged from 86.9 (SD 18.7) to 71.1 (SD 14.7) in individuals in the lowest (10) to the highest risk score category (25), and CKD prevalence ranged from 3.9 to 23.6%, respectively.

Analyses of Expression-Associated SNPs (eSNPs)

To obtain evidence for the presence of functional variants at the identified genomic risk loci and to prioritize genes in the associated regions, we focused on SNPs previously identified in genome-wide studies as significantly related to gene expression in liver ($n = 3,322$),¹¹ lymphocytes ($n = 29,094$)¹² or lymphoblastoid cell lines ($n = 10,823$).¹³ These expression SNPs (eSNPs) were then evaluated for their association with eGFRcrea, CKD, and eGFRcys from the discovery analysis. Table 4 shows that 9 of the 20 novel susceptibility loci identified for eGFRcrea (7/13 renal function loci) contained one or more significant eSNPs in one or more of the expression tissues queried. In addition, three of the previously identified loci (*SHROOM3*, *GATM*, and *CST3*) also contained at least one eSNP. The correlation (r^2) between the significant eSNP at each locus with the strongest LD to lead SNP ranged from 0.01 (*FBXO22*) to >0.9 (*DAB2*, *GCKR*; Table 4). The lead SNP in *GCKR* (rs1260326), a non-synonymous coding variant, was significantly associated with gene

expression of the neighboring *IFT172* gene. Further, the eSNP data supports *SLC7A9* as the important gene at the chromosome 19 susceptibility locus, as well as *ALMS1* at the susceptibility locus on chromosome 2p13, since an eSNP in perfect LD ($r^2=1$) with rs13538 in the HapMap CEU population is significantly associated with *ALMS1* transcript expression in lymphocytes. All eSNPs with significant association with at least one renal trait are listed in Supplementary Table 4.

Secondary Analyses: False Discovery Rate (FDR)

In order to further identify genomic loci for kidney function and disease, secondary analyses were conducted to identify SNPs that did not reach genome-wide significance but were associated with eGFR_{crea} or CKD at an FDR of 0.05 (p-value $<4.8 \times 10^{-6}$). After exclusion of all SNPs within 1 Mb of a genome-wide significant SNP, 9 additional loci for eGFR_{crea} were identified, 4 of which were also associated with eGFR_{cys} (Supplementary Table 5). We also compared the identified FDR-loci with the eSNP analysis; 3 regions of overlap were identified. The r^2 between each FDR SNP and the eSNP in highest LD with that FDR SNP ranged from 0.18 (*ARL15*) to 1.0 (*CASP9*, *CRKRS*), further supporting these genomic regions as loci of interest. Based on known biology, *PARD3B* and *CASP9* are particularly interesting genes emerging from FDR analyses. *PARD3B* is important in establishing cell polarity and localizes to tight junctions of epithelial cells;¹⁴ it is most expressed in fetal and adult kidney. *CASP9* is involved in the growth of metanephroi in the developing kidney.¹⁵

Discussion

Our principal findings are four-fold. We have identified 20 novel replicated loci in association with eGFR and CKD. Of these, 13 are likely to be involved in renal function and susceptibility to CKD, whereas 7 likely represent creatinine production or secretion loci. In aggregate, the 13 new renal function loci plus the three previously identified renal function loci account for 1.4% of the variation in eGFR_{crea}. We demonstrate altered transcript expression with SNPs at several of the identified loci, providing potential functional insight. Lastly, we provide suggestive evidence for an additional 9 eGFR_{crea}-associated loci using a false discovery rate metric, of which 4 loci are suspected to be related to renal function.

These findings extend previous knowledge of common genetic variation related to renal function indices. We have confirmed our prior findings, the identification of common risk variants at the *UMOD*, *SHROOM3*, and *STC1* loci as well as at two positive control loci (*GATM*, *CST*).¹⁰ We now highlight 13 novel loci not previously known to be associated with renal function in population-based studies. In the course of our work, we have also uncovered 7 loci likely influencing creatinine production or secretion. This underscores the importance of separating genetic loci that affect concentrations of a biomarker independent of underlying disease processes from those that truly reflect disease-association. Similar to what we have previously reported,¹⁰ we identified many more robust associations for eGFR_{crea} as compared to CKD. Nonetheless, nominal associations with CKD were identified for most of the renal function SNPs, showing that genetic variants associated with normal variation in eGFR_{crea} are also associated with the clinically important entity CKD.

Our findings highlight in several important ways how GWAS can aid in uncovering the genetic underpinnings of complex human traits and diseases as well as represent a first step towards a better understanding of physiological mechanisms and pathways. First, they provide novel information about the allelic architecture of known risk loci for genetic diseases of the kidney. Rare mutations in *SLC7A9*, *SLC34A1*, *ALMS1*, and *UMOD* are known to cause monogenetic diseases that feature a renal phenotype, underscoring the additional importance of common genetic susceptibility variants in these genes. This phenomenon is also observed for other complex traits and diseases; approximately 20% of

loci discovered in GWAS of a variety of complex traits are known to also harbor mutations that cause monogenic diseases.¹⁶

Second, our findings provide information about the genomic location of genetic variants associated with renal indices. Over 65% of the SNPs identified in our discovery analyses are located in or within 3.7 kb upstream of genes, and three of the variants are non-synonymous coding. This is in agreement with a recent study that reported an enrichment of trait-associated variants identified in GWAS at non-synonymous coding sites and a depletion of trait-associated variants in intergenic regions when compared to a random selection of variants on genotyping arrays.^{17;18}

Third, our replicated findings highlight the role of several pathways and mechanisms of importance in renal development and function. We identified common genetic variants in genes related to nephrogenesis (*ALMS1*, *VEGFA*, potentially *DACH1*), glomerular filtration barrier formation and podocyte function (*DAB2*, *PARD3B*, *VEGFA*), angiogenesis (*VEGFA*), solute transport (*SLC7A9*, *SLC34A1*), and metabolic functions of the kidney (*PRKAG2*, potentially *GCKR* and *LASS2*). Several of the genes we identified can be linked to the role of primary cilia (*ALMS1*, *GCKR/IFT172*, *PARD3B*); mutations in genes with a role in development and function of primary cilia are known to cause hereditary genetic diseases of the kidney such as polycystic kidney disease and nephronophthisis.¹⁹ The ability to uncover genetic variation in these genes in unselected individuals from population-based studies emphasizes their contribution to important mechanisms related to renal function in humans under physiological conditions and should provide interesting candidates for follow up in functional studies.

Finally, our data provide novel information about components of creatinine metabolism in humans and specifically about how creatinine is handled by epithelial cells of the proximal renal tubule. This may be of interest not only to physiologists but may also have consequences on the precision of GFR estimation from serum creatinine in the clinical setting.

Similar to what has been observed previously, we identified modest effects of the risk alleles on eGFR and CKD. Taken together, these renal function loci are associated with 1.4% of the variation in eGFR_{crea}. We observed substantial gradation of CKD prevalence across the genetic risk score, indicating the potential clinical relevance of these risk alleles. Most importantly, our findings point toward novel mechanisms that may lead to a better understanding of both renal development and the pathogenesis of CKD.

The strength of our analysis lies in the large sample size of 67,093 used for discovery, allowing us to uncover multiple loci despite the small effect size on eGFR and CKD. We restricted our analyses to population-based studies thereby avoiding potential bias from using case-control samples²⁰ or from potential counter-regulating disease processes. We employed several additional methods to enhance our ability to discover novel kidney disease susceptibility loci in this screen free of prior biological hypotheses including an FDR and eSNP approach. To enable discrimination of renal function loci from creatinine production and secretion loci, we used a separate complementary measure of glomerular filtration obtained from cystatin C.

Some limitations warrant mention. Our sample consists of white participants only, and it is uncertain whether these results would replicate in other ethnic groups. We used an indirect measure of GFR as estimated by the MDRD equation; gold-standard measures of glomerular filtration are not feasible in a large population-based setting. We used eGFR_{cys} to provide confirmatory evidence of which novel loci were indeed renal function loci. Loci that were not associated with eGFR_{cys} were designated as presumptive creatinine production or

secretion loci, but the smaller sample size with available cystatin C measures limits the power to confirm renal function loci. We may have falsely labeled some loci as unrelated to renal function based on an absent association with eGFR_{cys}, particularly *WDR37* and *WDR72*, which showed association of borderline significance with eGFR_{cys}. Lastly, several genes exist in the regions of interest. For many of the reported loci, we are unable to identify which is the most likely gene related to the SNP association based on statistical evidence, although we could address this question to some extent using the eSNP data.

Multiple common genetic variants are associated with indices of renal function and highlight the role of specific genes in nephrogenesis, podocyte function, angiogenesis, solute transport, and metabolic functions of the kidney.

Box 1. Genes of special interest at newly discovered susceptibility loci for kidney function and disease

SLC7A9

SLC7A9 encodes for an amino acid transporter in renal proximal tubule cells; mutations in *SLC7A9* cause cystinuria type B (OMIM #220100).²¹ Patients with cystinuria excrete elevated amounts of amino acids, resulting in the formation of stones in the urinary tract. *SLC7A9*-deficient mice display tubular and pelvic dilatation, tubular necrosis, and chronic interstitial nephritis.²²

SLC34A1

Mutations in *SLC34A1* cause hypophosphatemic nephrolithiasis/osteoporosis (OMIM #612286).²³ *SLC34A1* encodes the type IIa Na/Pi cotransporter, which is exclusively expressed in kidney and located in the brush border of renal proximal tubular cells, where it mediates reuptake of inorganic phosphate.²⁴

ALMS1

Mutations in *ALMS1* cause the autosomal recessive Almsstrom Syndrome (OMIM #203800), characterized by retinal degeneration, hearing loss, obesity, diabetes, and commonly renal insufficiency.^{25;26} Mutations in this gene are associated with age-dependent ciliopathies in the kidney.²⁷

DAB2

DAB2 is a cytoplasmatic adaptor protein expressed in renal proximal tubular cells,²⁸ where it is reported to represent the physical link between megalin and non-muscle myosin heavy polypeptide 9, encoded by *MYH9*.²⁹ Common variants in *MYH9* were recently identified as important susceptibility alleles for non-diabetic kidney disease in African Americans.^{30;31}

VEGFA

Encodes for vascular endothelial growth factor A, with roles in angiogenesis and vascular permeability.³² Renal podocytes produce large amounts of VEGFA, which is essential for glomerulogenesis and glomerular filtration barrier formation in animal models.³² *VEGFA* has been reported to affect ureteric bud growth during embryogenesis and hence may impact the number of nephrons.³³

GCKR

The product of *GCKR* inhibits hepatic glucokinase.³⁴ Common variants in *GCKR*, and specifically the missense SNP rs1260326 (P446L), are associated with a variety of human traits in genetic association studies, including serum triglycerides, fasting glucose, C-reactive protein, and uric acid as well as susceptibility to type 2 diabetes

(<http://www.genome.gov/GWastudies/>), highlighting the pleiotropy of this locus. eSNP analyses point to the role of the neighboring *IFT172* gene, which has a role in the formation of primary cilia.³⁵

PRKAG2

Rare *PRKAG2* variants cause a form of heart disease, featuring hypertrophic cardiomyopathy and the Wolff-Parkinson-White syndrome^{36;37} and sometimes enlarged kidneys.³⁸ Studies in transgenic mice indicate that these mutations cause a glycogen storage disease of the heart.³⁷ Several other hereditary glycogen storage diseases present with renal pathology such as renal tubular dysfunction.³⁹

DACH1

Dachshund homologue 1 is a transcription factor with a role in organogenesis; our data implicate a 100kb region within this gene. It is expressed in adult human kidney, as well as murine developing kidney, specifically glomerular podocytes and tubular epithelial cells.⁴⁰ *DACH1* may have a role in the development of the Mullerian duct.⁴¹ *DACH1* is part of the genetic network including *SIX* and *EYA*,⁴² mutations in which cause brachio-oto-renal syndrome.⁴³

LASS2

is highly expressed in the kidney and may be involved in cell growth.⁴⁴ A non-synonymous coding SNP in *LASS2*, rs267738 (E115A), was in perfect LD with the lead SNP in the region and of predicted damaging function.⁴⁵ *LASS2* has been implicated in the synthesis of specific ceramides.⁴⁶ Ceramides and their product sphingolipids are important in genetic diseases of the kidney,⁴⁷ and have a role in aging mechanisms.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Reference List

1. Coresh J, Selvin E, Stevens LA, Manzi J, Kusek JW, Eggers P, Van LF, Levey AS. Prevalence of chronic kidney disease in the United States. *JAMA* 2007;298:2038–2047. [PubMed: 17986697]
2. Meguid, ElNA.; Bello, AK. Chronic kidney disease: the global challenge. *Lancet* 2005;365:331–340. [PubMed: 15664230]
3. Go AS, Chertow GM, Fan D, McCulloch CE, Hsu CY. Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med* 2004;351:1296–1305. [PubMed: 15385656]
4. Smith DH, Gullion CM, Nichols G, Keith DS, Brown JB. Cost of medical care for chronic kidney disease and comorbidity among enrollees in a large HMO population. *J Am Soc Nephrol* 2004;15:1300–1306. [PubMed: 15100370]
5. Levey AS, Atkins R, Coresh J, Cohen EP, Collins AJ, Eckardt KU, Nahas ME, Jaber BL, Jadoul M, Levin A, Powe NR, Rossert J, Wheeler DC, Lameire N, Eknoyan G. Chronic kidney disease as a global public health problem: approaches and initiatives - a position statement from Kidney Disease Improving Global Outcomes. *Kidney Int* 2007;72:247–259. [PubMed: 17568785]
6. Fox CS, Larson MG, Leip EP, Culleton B, Wilson PW, Levy D. Predictors of new-onset kidney disease in a community-based population. *JAMA* 2004;291:844–850. [PubMed: 14970063]
7. Fox CS, Muntner P. Trends in diabetes, high cholesterol, and hypertension in chronic kidney disease among U.S. adults: 1988–1994 to 1999–2004. *Diabetes Care* 2008;31:1337–1342. [PubMed: 18436617]
8. Satko SG, Sedor JR, Iyengar SK, Freedman BI. Familial clustering of chronic kidney disease. *Semin Dial* 2007;20:229–236. [PubMed: 17555489]

9. Fox CS, Yang Q, Cupples LA, Guo CY, Larson MG, Leip EP, Wilson PW, Levy D. Genomewide linkage analysis to serum creatinine, GFR, and creatinine clearance in a community-based population: the Framingham Heart Study. *J Am Soc Nephrol* 2004;15:2457–2461. [PubMed: 15339995]
10. Kottgen A, Glazer NL, Dehghan A, Hwang SJ, Katz R, Li M, Yang Q, Gudnason V, Launer LJ, Harris TB, Smith AV, Arking DE, Astor BC, Boerwinkle E, Ehret GB, Ruczinski I, Scharpf RB, Ida Chen YD, de B I, Haritunians T, Lumley T, Sarnak M, Siscovick D, Benjamin EJ, Levy D, Upadhyay A, Aulchenko YS, Hofman A, Rivadeneira F, Uitterlinden AG, van Duijn CM, Chasman DI, Pare G, Ridker PM, Kao WH, Witteman JC, Coresh J, Shlipak MG, Fox CS. Multiple loci associated with indices of renal function and chronic kidney disease. *Nat Genet*. 2009
11. Schadt EE, Molony C, Chudin E, Hao K, Yang X, Lum PY, Kasarskis A, Zhang B, Wang S, Suver C, Zhu J, Millstein J, Sieberts S, Lamb J, GuhaThakurta D, Derry J, Storey JD, vila-Campillo I, Kruger MJ, Johnson JM, Rohl CA, van NA, Mehrabian M, Drake TA, Lusis AJ, Smith RC, Guengerich FP, Strom SC, Schuetz E, Rushmore TH, Ulrich R. Mapping the genetic architecture of gene expression in human liver. *PLoS Biol* 2008;6:e107. [PubMed: 18462017]
12. Goring HH, Curran JE, Johnson MP, Dyer TD, Charlesworth J, Cole SA, Jowett JB, Abraham LJ, Rainwater DL, Comuzzie AG, Mahaney MC, Almasy L, MacCluer JW, Kissebah AH, Collier GR, Moses EK, Blangero J. Discovery of expression QTLs using large-scale transcriptional profiling in human lymphocytes. *Nat Genet* 2007;39:1208–1216. [PubMed: 17873875]
13. Dixon AL, Liang L, Moffatt MF, Chen W, Heath S, Wong KC, Taylor J, Burnett E, Gut I, Farrall M, Lathrop GM, Abecasis GR, Cookson WO. A genome-wide association study of global gene expression. *Nat Genet* 2007;39:1202–1207. [PubMed: 17873877]
14. Kohjima M, Noda Y, Takeya R, Saito N, Takeuchi K, Sumimoto H. PAR3beta, a novel homologue of the cell polarity protein PAR3, localizes to tight junctions. *Biochem Biophys Res Commun* 2002;299:641–641. [PubMed: 12459187]
15. Hayashi M, Araki T. Caspase in renal development. *Nephrol Dial Transplant* 2002;17 Suppl 9:8–10. [PubMed: 12386274]
16. Hirschhorn JN. Genomewide association studies--illuminating biologic pathways. *N Engl J Med* 2009;360:1699–1701. [PubMed: 19369661]
17. Johnson AD, O'Donnell CJ. An open access database of genome-wide association results. *BMC Med Genet* 2009;10:6. [PubMed: 19161620]
18. Hindorf LA, Sethupathy P, Junkins HA, Ramos EM, Mehta JP, Collins FS, Manolio TA. Potential etiologic and functional implications of genome-wide association loci for human diseases and traits. *Proc Natl Acad Sci U S A* 2009;106:9362–9367. [PubMed: 19474294]
19. Davenport JR, Yoder BK. An incredible decade for the primary cilium: a look at a once-forgotten organelle. *Am J Physiol Renal Physiol* 2005;289:F1159–F1169. [PubMed: 16275743]
20. Monsees GM, Tamimi RM, Kraft P. Genome-wide association scans for secondary traits using case-control samples. *Genet Epidemiol*. 2009
21. Mattoo A, Goldfarb DS. Cystinuria. *Semin Nephrol* 2008;28:181–191. [PubMed: 18359399]
22. Feliubadalo L, Arbones ML, Manas S, Chillaron J, Visa J, Rodes M, Rousaud F, Zorzano A, Palacin M, Nunes V. Slc7a9-deficient mice develop cystinuria non-I and cystine urolithiasis. *Hum Mol Genet* 2003;12:2097–2108. [PubMed: 12915471]
23. Prie D, Huart V, Bakouh N, Planelles G, Dellis O, Gerard B, Hulin P, que-Blanchet F, Silve C, Grandchamp B, Friedlander G. Nephrolithiasis and osteoporosis associated with hypophosphatemia caused by mutations in the type 2a sodium-phosphate cotransporter. *N Engl J Med* 2002;347:983–991. [PubMed: 12324554]
24. Magagnin S, Werner A, Markovich D, Sorribas V, Stange G, Biber J, Murer H. Expression cloning of human and rat renal cortex Na/Pi cotransport. *Proc Natl Acad Sci U S A* 1993;90:5979–5983. [PubMed: 8327470]
25. Marshall JD, Hinman EG, Collin GB, Beck S, Cerqueira R, Maffei P, Milan G, Zhang W, Wilson DI, Hearn T, Tavares P, Vettor R, Veronese C, Martin M, So WV, Nishina PM, Naggert JK. Spectrum of ALMS1 variants and evaluation of genotype-phenotype correlations in Alstrom syndrome. *Hum Mutat* 2007;28:1114–1123. [PubMed: 17594715]

26. Joy T, Cao H, Black G, Malik R, Charlton-Menys V, Hegele RA, Durrington PN. Alstrom syndrome (OMIM 203800): a case report and literature review. *Orphanet J Rare Dis* 2007;2:49. [PubMed: 18154657]
27. Li G, Vega R, Nelms K, Gekakis N, Goodnow C, McNamara P, Wu H, Hong NA, Glynne R. A role for Alstrom syndrome protein, *alms1*, in kidney ciliogenesis and cellular quiescence. *PLoS Genet* 2007;3:e8. [PubMed: 17206865]
28. Nagai J, Christensen EI, Morris SM, Willnow TE, Cooper JA, Nielsen R. Mutually dependent localization of megalin and Dab2 in the renal proximal tubule. *Am J Physiol Renal Physiol* 2005;289:F569–F576. [PubMed: 15870384]
29. Hosaka K, Takeda T, Iino N, Hosojima M, Sato H, Kaseda R, Yamamoto K, Kobayashi A, Gejyo F, Saito A. Megalin and nonmuscle myosin heavy chain IIA interact with the adaptor protein Disabled-2 in proximal tubule cells. *Kidney Int* 2009;75:1308–1315. [PubMed: 19340093]
30. Kao WH, Klag MJ, Meoni LA, Reich D, Berthier-Schaad Y, Li M, Coresh J, Patterson N, Tandon A, Powe NR, Fink NE, Sadler JH, Weir MR, Abboud HE, Adler SG, Divers J, Iyengar SK, Freedman BI, Kimmel PL, Knowler WC, Kohn OF, Kramp K, Leehey DJ, Nicholas SB, Pahl MV, Schelling JR, Sedor JR, Thornley-Brown D, Winkler CA, Smith MW, Parekh RS. MYH9 is associated with nondiabetic end-stage renal disease in African Americans. *Nat Genet* 2008;40:1185–1192. [PubMed: 18794854]
31. Kopp JB, Smith MW, Nelson GW, Johnson RC, Freedman BI, Bowden DW, Oleksyk T, McKenzie LM, Kajiyama H, Ahuja TS, Berns JS, Briggs W, Cho ME, Dart RA, Kimmel PL, Korbet SM, Michel DM, Mokrzycki MH, Schelling JR, Simon E, Trachtman H, Vlahov D, Winkler CA. MYH9 is a major-effect risk gene for focal segmental glomerulosclerosis. *Nat Genet* 2008;40:1175–1184. [PubMed: 18794856]
32. Eremina V, Baelde HJ, Quaggin SE. Role of the VEGF--a signaling pathway in the glomerulus: evidence for crosstalk between components of the glomerular filtration barrier. *Nephron Physiol* 2007;106:32–37.
33. Karihaloo A, Karumanchi SA, Cantley WL, Venkatesha S, Cantley LG, Kale S. Vascular endothelial growth factor induces branching morphogenesis/tubulogenesis in renal epithelial cells in a neuropilin-dependent fashion. *Mol Cell Biol* 2005;25:7441–7448. [PubMed: 16107693]
34. Iynedjian PB. Molecular physiology of mammalian glucokinase. *Cell Mol Life Sci* 2009;66:27–42. [PubMed: 18726182]
35. Gorivodsky M, Mukhopadhyay M, Wilsch-Braeuninger M, Phillips M, Teufel A, Kim C, Malik N, Huttner W, Westphal H. Intraflagellar transport protein 172 is essential for primary cilia formation and plays a vital role in patterning the mammalian brain. *Dev Biol* 2009;325:24–32. [PubMed: 18930042]
36. Blair E, Redwood C, Ashrafian H, Oliveira M, Broxholme J, Kerr B, Salmon A, Ostman-Smith I, Watkins H. Mutations in the gamma(2) subunit of AMP-activated protein kinase cause familial hypertrophic cardiomyopathy: evidence for the central role of energy compromise in disease pathogenesis. *Hum Mol Genet* 2001;10:1215–1220. [PubMed: 11371514]
37. Arad M, Benson DW, Perez-Atayde AR, McKenna WJ, Sparks EA, Kanter RJ, McGarry K, Seidman JG, Seidman CE. Constitutively active AMP kinase mutations cause glycogen storage disease mimicking hypertrophic cardiomyopathy. *J Clin Invest* 2002;109:357–362. [PubMed: 11827995]
38. Burwinkel B, Scott JW, Buhrer C, van Landeghem FK, Cox GF, Wilson CJ, Grahame HD, Kilimann MW. Fatal congenital heart glycogenosis caused by a recurrent activating R531Q mutation in the gamma 2-subunit of AMP-activated protein kinase (PRKAG2), not by phosphorylase kinase deficiency. *Am J Hum Genet* 2005;76:1034–1049. [PubMed: 15877279]
39. Ozen H. Glycogen storage diseases: new perspectives. *World J Gastroenterol* 2007;13:2541–2553. [PubMed: 17552001]
40. Ayres JA, Shum L, Akarsu AN, Dashner R, Takahashi K, Ikura T, Slavkin HC, Nuckolls GH. DACH: genomic characterization, evaluation as a candidate for postaxial polydactyly type A2, and developmental expression pattern of the mouse homologue. *Genomics* 2001;77:18–26. [PubMed: 11543628]
41. Davis RJ, Harding M, Moayed Y, Mardon G. Mouse Dach1 and Dach2 are redundantly required for Mullerian duct development. *Genesis* 2008;46:205–213. [PubMed: 18395837]

42. Ikeda K, Watanabe Y, Ohto H, Kawakami K. Molecular interaction and synergistic activation of a promoter by Six, Eya, and Dach proteins mediated through CREB binding protein. *Mol Cell Biol* 2002;22:6759–6766. [PubMed: 12215533]
43. Abdelhak S, Kalatzis V, Heilig R, Compain S, Samson D, Vincent C, Weil D, Cruaud C, Sahly I, Leibovici M, Bitner-Glindzicz M, Francis M, Lacombe D, Vigneron J, Charachon R, Boven K, Bedbeter P, Van RN, Weissenbach J, Petit C. A human homologue of the *Drosophila* eyes absent gene underlies branchio-oto-renal (BOR) syndrome and identifies a novel gene family. *Nat Genet* 1997;15:157–164. [PubMed: 9020840]
44. Pan H, Qin WX, Huo KK, Wan DF, Yu Y, Xu ZG, Hu QD, Gu KT, Zhou XM, Jiang HQ, Zhang PP, Huang Y, Li YY, Gu JR. Cloning, mapping, and characterization of a human homologue of the yeast longevity assurance gene LAG1. *Genomics* 2001;77:58–64. [PubMed: 11543633]
45. Ng PC, Henikoff S. Predicting deleterious amino acid substitutions. *Genome Res* 2001;11:863–874. [PubMed: 11337480]
46. Mizutani Y, Kihara A, Igarashi Y. Mammalian Lass6 and its related family members regulate synthesis of specific ceramides. *Biochem J* 2005;390:263–271. [PubMed: 15823095]
47. Shayman JA. Sphingolipids: their role in intracellular signaling and renal growth. *J Am Soc Nephrol* 1996;7:171–182. [PubMed: 8785385]




Figure 1. Genome-wide $-\log_{10}$ p-value plot from Stage 1: Discovery analysis of eGFRcrea (**A**), CKD (**B**) and eGFRcys (**C**).




Figure 2.




Figure 3.

Table 1
Study Sample Characteristics in Stage 1 Discovery and Stage 2 Replication samples, CKDGen Consortium

Study	Sample Size eGFR _{crea} /CKD/eGFR _{cys}	Women %	Age (years)	eGFR _{crea} (ml/min/1.73 m ²)	eGFR _{cys} (ml/min/1.73 m ²)	CKD* %
Stage 1: Discovery						
AGES	3219/3219/NA	58.0	76.4 (5.5)	73.0 (20.0)	NA	24.3
Amish	1211/NA/783	49.0	49.3 (16.8)	93.8 (19.6)	114.9 (17.9)	2.9
ARIC	8069/8069/6430	53.0	62.0 (6.1)	81.3 (17.7)	84.1 (19.7)	9.1
ASPS	850/850/NA	56.8	65.2 (8.1)	96.5 (39.9)	NA	8.1
BLSA	723/723/NA	46.1	70.4 (15.2)	80.3 (23.1)	NA	17.4
CHS	3259/3259/2820	53.0	72.3 (5.4)	80.0 (22.6)	79.9 (18.3)	18.5
ERF	2079/2079/NA	56.3	49.2 (14.0)	93.5 (21.4)	NA	3.8
Family Heart Study	883/883/NA	51.1	55.5 (11.1)	88.5 (19.4)	NA	4.4
Framingham Heart Study	7782/4140/2992	54.3	51.2 (14.0)	92.1 (21.7)	77.9 (16.9)	10.7
KORA F3	1641/1641/1642	50.5	62.5 (10.1)	78.7 (19.0)	111.9 (26.6)	10.7
KORA F4	1814/1814/1811	51.3	60.9 (8.9)	85.1 (20.2)	109.7 (26.2)	7.0
Korcula	888/888/NA	64.0	56.3 (13.9)	87.3 (20.6)	NA	7.5
Micros	1201/1201/1248	56.8	46.2 (16.1)	94.6 (20.9)	107.4 (23.6)	3.8
ORCADES	704/704/NA	53.6	54.1 (15.2)	89.4 (20.7)	NA	6.8
Rotterdam Study I	4390/4390/NA	61.4	70.0 (9.0)	77.1 (17.2)	NA	13.7
Rotterdam Study II	1863/1863/NA	54.5	64.8 (8.0)	81.3 (17.2)	NA	9.1
NHSPS	565/565/NA	53.1	51.7 (18.3)	90.9 (22.1)	NA	5.7
SHIP	3231/3228/3231	51.7	54.5 (15.3)	90.5 (23.6)	97.1 (25.31)	7.7
Vis	768/768/NA	58.6	56.9 (15.2)	88.1 (22.1)	NA	6.9
WGHS	21953/21953/NA	100.0	54.7 (7.1)	90.3 (22.5)	NA	6.1
Total	67093/62237/20957					
Stage 2: Replication						
ARIC	944/944/751	54.8	61.9 (6.13)	81.6 (16.2)	85.4 (19.6)	7.5
GENOA	1056/1056/NA	56.2	59.0 (10.2)	85.2 (22.8)	NA	12.0
Family Heart Study	1537/1537/NA	57.1	47.8 (13.3)	93.7 (19.7)	NA	2.2
Gutenberg Heart Study	3180/3180/NA	48.7	55.9 (10.9)	87.1 (16.6)	NA	3.8

Study	Sample Size eGFRcrea/CKD/eGFRcys	Women %	Age (years)	eGFRcrea (ml/min/1.73 m ²)	eGFRcys (ml/min/1.73 m ²)	CKD* %
Stage 1: Discovery						
Health ABC	1663/1663/1663	47.2	73.7 (2.9)	70.1 (14.5)	77.3 (20.1)	25.1
HPFS	818/818/NA	0	64.7 (8.3)	85.2 (22.7)	NA	9.5
KORA F3	1498/1498/1498	52.5	51.6 (13.3)	95.2 (24.2)	123.5 (29.5)	4.0
KORA F4	1202/1202/1202	52.3	49.2 (15.4)	92.9 (23.2)	118.4 (27.5)	5.9
Nurses Health Study	786/786/NA	100.0	59.5 (6.5)	86.2 (22.1)	NA	10.7
POPGEN	1163/1163/NA	40.6	53.8 (14.4)	88.1 (18.8)	NA	5.1
SAPALDIA	6031/6031/NA	50.2	52.2 (11.4)	90.7 (17.3)	NA	2.9
SAPHIR	1733/1733/NA	37.0	51.4 (6.0)	91.7 (16.0)	NA	1.1
SORBS	892/892/NA	58.3	48.5 (15.8)	92.6 (19.3)	NA	4.1
SPLIT	479/NA/NA	58.9	49.2 (14.5)	95.59 (24.1)	NA	3.5
Total	22982/22503/5114					

* Total of 5807 CKD cases in the Stage 1 Discovery sample and 1366 CKD cases in Stage 2 Replication.

Abbreviations: eGFRcrea: estimated glomerular filtration rate by serum creatinine, eGFRcys: estimated glomerular filtration rate by serum cystatin C, CKD: chronic kidney disease, NA: not available.

Table 2
Genome-wide Significant Loci: SNP Association with Renal Traits in Stage 1 Discovery And Stage 2 Replication Meta-Analyses

Trait	SNP ID	Chr	position (b36)	Genes Within 60 kb	Known Loci	SNP function	Minor Allele Freq#	Trait Discovery P-value*	Trait Replication P-value	Trait Meta- analysis P-value	eGFRcys P-value§
eGFRcrea*	rs17319721	4	77587871	SHROOM3,FLJ25770		intronic	0.43	1.1E-19	NA	NA	NA
eGFRcrea	rs10109414	8	23807096	STC1		intergenic	0.42	1.0E-08	NA	NA	NA
eGFRcrea	rs2453533	15	43428517	GATM,SPATA5L1		intergenic	0.38	4.6E-22	NA	NA	NA
eGFRcrea*	rs12917707	16	20275191	UMOD,FLJ20581,GP2,PDILT		upstream	0.18	1.2E-20	NA	NA	NA
eGFRcys	rs911119	20	23560737	CST3,CST4,CST9		intergenic	0.21	2.3E-138	NA	NA	NA
Novel Renal Function Loci											
eGFRcrea	rs1933182	1	109801361	SYPL2,ATXN7L2,CYB561D1,PSMA5,AMIGO1,SORT1		intergenic	0.30	1.3E-08	1.9E-01	1.2E-07	5.0E-02
eGFRcrea	rs267734	1	149218101	ANXA9,FAM63A,PRUNE,BNIP1,LASS2,SETDB1		upstream	0.20	5.2E-09	1.1E-04	1.2E-12	9.7E-03
CKD	rs16864170	2	5825331	SOX11		intergenic	0.05	4.5E-08	3.5E-01	1.6E-07	2.9E-02
eGFRcrea	rs1260326	2	27584444	GCKR,IFT172,FNDCA		non-synonymous	0.41	1.3E-10	1.1E-04	3.0E-14	3.7E-03
eGFRcrea	rs13538	2	73721836	NAT8,NAT8B,ALMS1		non-synonymous	0.23	2.6E-08	7.2E-07	4.5E-14	1.6E-04
eGFRcrea	rs347685	3	143289827	TFDP2		unknown	0.28	7.0E-09	1.4E-03	3.0E-11	1.2E-03
eGFRcrea	rs11959928	5	39432889	DAB2,C9		intronic	0.44	1.8E-11	5.6E-07	1.4E-17	1.6E-04
eGFRcrea	rs6420094	5	176750242	SLC34A1,GRK6,RGS14,LMAN2,PRR7,F12,PFN3		intronic	0.34	3.8E-12	6.6E-04	1.0E-14	1.3E-05
eGFRcrea	rs881858	6	43914587	VEGFA		intergenic	0.28	2.2E-11	7.7E-04	9.4E-14	3.7E-03
CKD*	rs7805747	7	151038734	PRKAG2		intronic	0.24	8.6E-09	7.7E-05	4.2E-12	8.1E-07
eGFRcrea	rs4744712	9	70624527	PIP5K1B,FAM122A		intronic	0.39	7.2E-10	6.6E-05	8.3E-14	6.8E-03
eGFRcrea	rs4014195	11	65263398	RNASEH2C,DKFZp761E19.8,HTATIP,OVOL1		intergenic	0.35	3.3E-08	1.4E-01	1.4E-07	2.7E-02
eGFRcys	rs653178	12	110492139	ATXN2		intronic	0.50	3.8E-08	1.4E-04	3.5E-11	3.5E-11
eGFRcrea	rs626277	13	71245697	DACH1		intronic	0.40	2.9E-10	1.0E-02	2.5E-11	6.5E-04
eGFRcrea	rs1394125	15	73946038	UBE2Q2,FBXO22		intronic	0.35	3.7E-10	4.7E-08	3.3E-17	6.9E-04
eGFRcrea	rs12460876	19	38048731	SLC7A9,CCDC123,ECAT8		intronic	0.39	5.5E-09	2.5E-07	3.2E-15	1.2E-02
Novel Creatinine Production and Secretion Loci											
eGFRcrea	rs7422339**	2	211248752	CPS1		non-synonymous	0.32	2.4E-09	2.6E-07	1.2E-15	1.9E-01
eGFRcrea	rs2279463	6	160588379	SLC22A2		intronic	0.12	8.7E-10	1.7E-03	5.5E-12	4.8E-01

Trait	SNP ID	Chr	position (b36)	Genes Within 60 kb	SNP function	Minor Allele Freq#	Trait Discovery P-value*	Trait Replication P-value	Trait Meta- analysis P-value	eGFRcys P-value§
eGFRcrea	rs6465825	7	77254375	TMEM60;RSBN1L;PHTF2	intergenic	0.39	3.5E-09	3.5E-02	1.5E-09	7.4E-01
eGFRcrea	rs10794720	10	1146165	WDR37	intronic	0.08	2.1E-08	4.7E-02	1.2E-08	8.9E-02
eGFRcrea	rs10774021	12	219559	SLC6A13 ;JARID1A;SLC6A12	intronic	0.36	6.7E-09	7.1E-02	1.4E-09	7.9E-01
eGFRcrea	rs491567	15	51733885	WDR72	intronic	0.22	1.3E-08	1.0E-05	2.7E-13	8.1E-02
eGFRcrea	rs9895661	17	56811371	BCAS3 ;TBX2;C17orf82	intronic	0.19	1.4E-08	3.0E-08	1.1E-15	2.8E-01

* Only rs12917707 at the UMOD locus demonstrated significant heterogeneity.

The minor allele based on sample size weighted mean allele frequency in the discovery cohorts is modeled.

* SNP also significantly associated with at least one other kidney trait at $p < 5 \times 10^{-8}$.

** SNP rs7422339 has changed the SNP identifier in the latest dbSNP release,current name is rs1047891.

§ Results of eGFRcys from Discovery and Replication combined. For ATXN2, the p-value for eGFRcrea from Stage 1 + Stage 2 is 0.0005. NA: known loci were not replicated in additional study samples. Genes within 60 kb were based on RefSeq genes (b36). The gene closest to the SNP is listed first and bold if the SNP is located within the gene. Other genes in the region are listed after ",".

Abbreviations: Chr=chromosome.

Table 3
Association of Replicated Renal Function Susceptibility Loci with CKD

Chr	position (b36)	Genes Nearby	modeled allele [#]	OR CKD [§]	95% CI	p-value
rs267734	1 149218101	ANXA9,FAM63A,PRUNE,BNIP1,LASS2,SETDB1	c	0.93	0.88–0.97	2.6E-03
rs1260326	2 27584444	GCKR ,IFT172,FNDC4	t	0.96	0.93–1.00	8.7E-02
rs13538	2 73721836	NAT8 ,NAT8B,ALMS1	g	0.93	0.89–0.98	6.3E-03
rs347685	3 143289827	TFDP2	c	0.93	0.89–0.97	1.2E-03
rs11959928	5 39432889	DAB2 ,C9	a	1.08	1.04–1.12	1.5E-05
rs6420094	5 176750242	SLC34A1 ,GRK6,RGS14,LMAN2,PRR7,F12,PFN3	g	1.08	1.03–1.12	2.1E-04
rs881858	6 43914587	VEGFA	g	0.93	0.89–0.97	3.7E-03
rs7805747	7 151038734	PRKAG2	a	1.19	1.13–1.25	4.2E-12
rs4744712	9 70624527	PIP5K1B ,FAM122A	a	1.06	1.02–1.10	7.0E-04
rs653178	12 110492139	ATXN2	t	0.96	0.92–1.00	5.8E-02
rs626277	13 71245697	DACHI	c	0.94	0.91–0.98	4.7E-03
rs1394125	15 73946038	UBE2Q2 ,FBXO22	a	1.08	1.04–1.13	8.0E-05
rs12460876	19 38048731	SLC7A9 ,CCDC123,ECAT8	c	0.93	0.89–0.96	2.6E-04

[#]The minor allele based on sample size weighted mean allele frequency in the discovery cohorts is modeled.

[§]Results based on Stage 1: Discovery and Stage 2: Replication combined. Genes within 60 kb were based on RefSeq genes. The gene closest to the SNP is listed first and bold if the SNP is located within the gene. Other genes in the region are listed after “,”. P-values from the discovery screen are corrected for genomic control before and after meta-analysis.

Abbreviations: OR: odds ratio, CKD: chronic kidney disease, Chr=chromosome.

Table 4

Significant eSNPs associated with eGFRcrea, CKD, or eGFRcys

P-values in bold are $<1/n$, with n being the number of eSNPs queried (3322 for liver, 10823 for lymphoblastoid cell lines (LCL), and 29094 for lymphocytes).

SNP	Tissue	chr	Genes within 60kb of SNP	p-value eSNP	expressed gene	p-value eGFRcys	p-value CKD	p-value eGFRcrea	r ² to top SNP	top SNP
rs1260326	lymphocytes, LCL *	2	GCKR;IFT172, FNDC4	7.0E-12	IFT172 [#]	6.4E-03	1.8E-01	1.3E-10	identical	rs1260326
rs10198549	lymphocytes, liver *	2	ALMS1	4.8E-09	ALMS1 [#]	9.1E-04	1.4E-01	3.8E-07	1	rs13538
rs6440052	lymphocytes	3	TFDP2;ATP1B3	5.2E-22	ATP1B3	1.5E-02	3.0E-04	5.0E-07	0.67	rs347685
rs4256249	liver	4	SHROOM3	4.3E-06	SHROOM3	2.7E-01	8.0E-02	1.0E-05	0.06	rs17319721
rs335223	lymphocytes	5	DAB2;C9	4.7E-04	DAB2	5.5E-03	2.2E-05	1.7E-10	0.93	rs11959928
rs1544457	LCL	7	TMEM60;RSBN1L,PHTF2	2.9E-08	PTPN12 [#]	2.9E-01	1.8E-01	4.4E-07	0.72	rs6465825
rs409783	liver	8	STC1	5.5E-07	C2orf29	1.3E-02	4.2E-01	3.1E-04	0.29	rs10109414
rs7035163	liver	9	FAM122A;PIP5K1B	2.4E-11	FAM122A	8.0E-01	2.4E-02	4.0E-05	0.19	rs4744712
rs11062357	lymphocytes	12	JARID1A;SLC6A13	5.5E-07	JARID1A	5.3E-01	1.9E-01	2.4E-06	0.37	rs10774021
rs335675	lymphocytes, liver *	15	FBXO22;NRG4, UBE2Q2	2.1E-03	FBXO22 [#]	2.0E-04	4.3E-01	9.4E-08	0.01	rs1394125
rs16967572	liver	19	CCDC123;RHPN2, SLC7A9,C19orf40	9.6E-13	SLC7A9	5.3E-02	1.9E-01	7.2E-05	0.58	rs12460876

* Other significant eSNPs in the same region were significantly associated with kidney traits in the second tissue listed. The SNPs, expressed genes, and the r^2 to the lead SNP in the region can be found in Supplementary Table 6.

Other significant eSNPs in the extended gene region were significantly associated with eGFRcrea and expression of other genes nearby. R^2 to the lead SNP in the region was lower. The SNPs, expressed genes, and r^2 to the lead SNP are listed in Supplementary Table 6. Additionally, eSNPs in creatinine and cystatin production loci GATM and CST3 were significantly associated with eGFRcrea and eGFRcys, respectively, in lymphocytes and are listed in Supplementary Table 4. Genes within 60 kb were based on RefSeq genes. The gene closest to the SNP is listed first and separated by other genes in the region by ";".